

# Disinfecting hygiene powder



## Documented quality hygiene

The scientific formulated powder is safe and easy to use and is designed to improve the animals surrounding environment by drying out and repressing noxious pathogens. Basic® hygiene powder reduces moisture and potentially noxious gases like ammonia. Use Basic® hygiene powder regularly to prevent disease, and to ensure a better environment for healthier animals.

- High quality at discount price
- Basic® has a documented effect
- Represses hatching conditions for flies
- Represses the growth of microorganisms
- Good absorption – strong drying out effect
- Does not lump in storage
- Good binder of ammonia
- Good for all kinds of animals and housing systems
- Improves the housing environment
- Reduces the smell of ammonia
- Freshens up the barn

## Description:

Basic® hygiene powder is composed of carefully selected materials, including minerals, plant components and a disinfectant active substance. Basic® hygiene powder is used to improve sanitary conditions for all kinds of livestock and in all housing systems. Basic® can be applied manually, with machines or be mixed with other bedding before application.

## Application areas:

Basic® hygiene powder is used for all kinds of animals and housing systems to dry out, decrease the pressure of infection and to improve the well-being of the animals.



### Basic® has proven effective against:

- E-coli
- Salmonella Thyphimorium
- Streptococcus Suis Serovar 7
- Streptococcus Uberis
- Staphylococcus Aureus
- Klebsiella Pneumoniae
- Aspergillus Niger
- Clostridium Perfringens
- Campylobacter Jejuni
- Dishelobactor Nodosus



[www.destek.dk](http://www.destek.dk)

Destek ApS

Orebovej 29

DK-4293 Dianalund

Phone: +45 3313 1125

Mail: [info@destek.dk](mailto:info@destek.dk)



# Disinfecting hygiene powder



## Instructions for use

### Basic® hygiene powder - laying hens

Apply Basic® in an even layer in the nests, according to 50-60 g per nest (a handful). If Basic® is used on the floor, it should be applied with 50-60 g per m<sup>2</sup> floor area.

After moving in a new group of hens, Basic® should be used daily the first 4 days and afterwards regularly 2-3 times a week. Basic® can beneficially be applied on the floor before bedding, such as straw or wood shavings, is spread. But Basic® can also be mixed with the common bedding before application.

By consequent use of Basic® hygiene powder at least 2-3 times a week, a dry environment is ensured, ammonia is bound and moist is absorbed. This way it is avoided that the bedding becomes too moist, because a moist bedding can lead to ammonia, which can result in etching and damages to the foot pads.

### Basic® hygiene powder - broiler chickens

After all chickens are removed and the barn is cleaned thoroughly, Basic® can beneficially be applied in the empty barn in an even layer. This is typically done with a spreading trolley, where Basic® is used according to 50-60 g per m<sup>2</sup> floor area.

After that the common bedding such as straw or wood shavings is applied, before Basic® is spread again on top of the common bedding according to 50-60 g per m<sup>2</sup> floor area. Then the new chickens are moved in and Basic® is applied 1-2 times a week until slaughter, according to 50-60 g per m<sup>2</sup> floor area.

In case of problems like a higher pressure of infection, or under especially moist conditions, the consumption of Basic® should be increased according to 75-100 g per m<sup>2</sup> floor area. The usage of Basic® should continue at this level, until the barn level is back to normal again.



#### Basic® has proven effective against:

- E-coli
- Salmonella Thyphimorium
- Streptococcus Suis Sero var 7
- Streptococcus Uberis
- Staphylococcus Aureus
- Klebsiella Pneumoniae
- Aspergillus Niger
- Clostridium Perfringens
- Camphylobacter Jejuni
- Dishelobactor Nodosus



[www.destek.dk](http://www.destek.dk)

Destek ApS

Orebovej 29

DK-4293 Dianalund

Phone: +45 3313 1125

Mail: [info@destek.dk](mailto:info@destek.dk)

